

ILLINOIS
NATURAL
HISTORY
SURVEY

Black-crowned Night-Herons of the Lake Calumet Region, Chicago, Illinois



Article 3

Nesting Ecology of Black-crowned Night-Herons at Lake Calumet Wetlands

Jeffrey M. Levensgood, Walter J. Marcisz,
Allison M. Klement, and Margaret A. Kurcz

Article 4

Population Trends in a Black-crowned Night-Heron Colony at Lake Calumet Wetlands

Walter J. Marcisz, Jeffrey M. Levensgood,
Allison M. Klement, and Margaret A. Kurcz

Illinois Natural History Survey, David L. Thomas, Chief
A Division of the Illinois Department of Natural Resources

Illinois Natural History Survey
Distribution Office
607 E. Peabody Dr.
Champaign, IL 61820

Citations:

Levengood, J.M., W.J. Marcisz, A.M. Klement, and M.A. Kurcz. 2005. Nesting ecology of Black-crowned Night-Herons at Lake Calumet Wetlands. *Illinois Natural History Survey Bulletin* 37(3):95–108.

Marcisz, W.J., J.M. Levengood, A.M. Klement, and M.A. Kurcz. 2005. Population trends in a Black-crowned Night-Heron colony at Lake Calumet Wetlands. *Illinois Natural History Survey Bulletin* 37(4):109–118.

Editor: Charles Warwick

Photos on frontis: Michael Jeffords and Maggie Kurcz, Illinois Natural History Survey and Marv Piwoni, Illinois Waste Management Research Center

US ISSN 0073-4918


Printed by authority of the State of Illinois
P0076599—.75M—08-05

Printed with soy ink on recycled and recyclable paper.

Equal opportunity to participate in programs of the Illinois Department of Natural Resources (IDNR) and those funded by the U.S. Fish and Wildlife Service and other agencies is available to all individuals regardless of race, sex, national origin, disability, age, religion, or other non-merit factors. If you believe you have been discriminated against, contact the funding source's civil rights office and/or the Equal Employment Opportunity Officer, IDNR, One Natural Resources Way, Springfield, IL 62702-1271; 217/785-0067; TTY 217/782-9175.


ILLINOIS
NATURAL
HISTORY
SURVEY

Black-crowned Night-Herons of the Lake Calumet Region, Chicago, Illinois



ILLINOIS
NATURAL
HISTORY
SURVEY

Black-crowned Night-Herons of the Lake Calumet Region, Chicago, Illinois



Article 3

Nesting Ecology of Black-crowned Night-Herons at Lake Calumet Wetlands

Jeffrey M. Levensgood, Walter J. Marcisz,
Allison M. Klement, and Margaret A. Kurcz

Correspondence to:

Jeffrey M. Levensgood
Center for Wildlife and Plant Ecology
Illinois Natural History Survey
607 E. Peabody Dr.
Champaign IL 61820
217/ 333-6767
jleven@inhs.uiuc.edu

CONTENTS

ABSTRACT	95
INTRODUCTION	95
METHODS	96
2002	96
2003	96
RESULTS	96
Nesting Habitat at Lake Calumet Wetlands 1984–2003	96
Arrival and Courtship	98
Nesting Phenology	98
Clutch Size	101
Nest, Egg, and Nesting Fate	101
Fledging and Dispersal	103
Productivity	103
DISCUSSION	106
LITERATURE CITED	107
ACKNOWLEDGEMENTS	108

ABSTRACT

We examined the nesting ecology of a Black-crowned Night-Heron (*Nycticorax nycticorax*) colony located at wetlands associated with Lake Calumet in south Cook County, Illinois, during the 2002 and 2003 nesting seasons. This area of southeastern Chicago has been greatly impacted by heavy industry, solid and chemical waste disposal, urbanization, and altered hydrology.

Black-crowned Night-Herons (BCNH) have nested at five known locations at Lake Calumet wetlands during 1984–2003. Emergent cover (giant reed, *Phragmites australis*) was of primary importance to this colony for nesting during that time. Cottonwoods (*Populus deltoides*) also were used for nesting from the late 1980s to mid-1990s.

During 1993–2003 the herons began arriving at the colony as early as March 10. During the two years of this study the earliest indications of nest building and courtship occurred during the first week in April; the first pairs and precopulatory displays were observed during second week of April in both years. The egg-laying period extended from April 20 to June 12 in 2002, and from April 16 to May 27/28 in 2003. Hatching occurred from mid-May to the first few days of July in 2002, and from mid-May to June 19/20 in 2003. Juvenile dispersal in 2002 occurred from mid-July through late August, and from early July through mid-August in 2003.

Reproductive parameters in BCNH nesting at the north end of Indian Ridge Marsh (IRM), the primary nesting location for this colony in both years, were typical for this species. In 2002 the “recruitment” rate (number of young/nest surviving to 15 days) of 1.74 young/pair was below the threshold of 2.0–2.1 young/nesting pair thought to be necessary to maintain BCNH populations. However, recruitment increased to 2.22 young/pair in 2003, which was among the highest previously reported. The most important cause of nest failure was poorly constructed (flat) nests which allowed the eggs to roll out into the water. Although some eggs were lost to gulls and some hatched young were taken by unknown mammalian or avian predators, predation was not an important cause of nest losses at IRM.

INTRODUCTION

The Calumet region of southwestern Lake Michigan was once a vast complex of glacial lakes, wetlands, and sand prairies. This region is now one of most heavily-industrialized in the U.S. and has been greatly impacted through industrial activities, waste disposal and discharge, urbanization, and changes to surface and groundwater hydrology. In spite of extensive habitat loss and degradation, the area remains among the most biologically diverse in the state of Illinois.

A number of wetland-dependent breeding birds of concern nest in marshes adjacent to Lake Calumet, including the state-endangered Black-crowned Night-Heron (*Nycticorax nycticorax*). One of the largest remaining breeding colonies in Illinois, this population is of considerable interest to resource professionals, environmental groups, and the conservation-minded public. Although BCNH populations have increased nationwide since banning of DDT and other persistent pesticides, Illinois has not enjoyed this recovery.

Young-of-the-year BCNH were reported in the Calumet marshes as early as 1874 (Nelson 1876–1877). Throughout much of the early part of the 20th century the Lake Calumet BCNH colony was located along the Calumet River just north of the confluence with the Grand Calumet River (Landing 1986). The BCNH have nested at wetlands adjacent to Lake Calumet since the Thomas J. O’Brien Lock and Dam went into operation in the late 1960s. The number of BCNH nesting at these wetlands has fluctuated widely over the last two decades. However, this population remained relatively stable at between 300 and 400 pairs during 1997–2003 (Marcisz et al. 2005).

Interest in the rehabilitation of these wetlands as part of the Calumet Open Space Reserve created the need for a better understanding of the ecology of this urban BCNH colony, in order to aid conservation planning and guide management activities. Accordingly, we examined the nesting ecology of BCNH at Lake Calumet’s IRM during the 2002 and 2003 seasons.

METHODS

2002

The activities of BCNH were observed for two-to-four-hour periods on eight occasions during April 5 to May 10 to document the timing of breeding activities. Nest building, pairing, breeding, and nesting behaviors were recorded. Nest monitoring was initiated at IRM on May 15, when 30 nests were marked by placing plastic flagging on *Phragmites* stems adjacent to each nest. The number of eggs and chicks was recorded for active nests. Additional nests were added May 20 (n= 9), May 26 (n= 6), May 30 (n= 2), and June 4 (n= 1). Nests were checked every four to seven days depending on weather and staff availability. The number and disposition of eggs (cracking, pipping, hatched, fell out, depredated, missing) and nestlings (present, missing, depredated) were recorded during each visit; eggs were numbered with a nontoxic marker. Twelve nests located in cottonwood trees at a colony located 13 km to the east at ISPAT Inland Steel (IIS), East Chicago, IN, were monitored for comparison; each monitored nest was numbered on a hand-drawn map of the site and observed from an elevated vantage point.

We conducted post-breeding surveys at IRM every two weeks during June–August 2002 to monitor the relative abundance of juvenile BCNH from about the time the first young became flighted through dispersal. These surveys consisted of walking an established transect adjacent to and the length of the colony at a slow pace, counting and aging the herons as the observer passed by them. To avoid double-counting, herons that took flight at the observer's approach and landed within an unsurveyed portion of the colony were not counted at that time.

2003

The activities of BCNH were observed for 1- to 2.5-hour periods on 11 occasions during April 9 to May 13. Nest building, pairing, breeding, and nesting behaviors were recorded. Nest monitoring at IRM was initiated on May 16; a total of 55 nests were marked (n= 20 on 5/16, 23 on 5/21, and 12 on 5/29). A total of 17 nests located at IIS were also monitored. Nest monitoring at IRM and IIS, respectively, was concluded July 10 and 23. Dates of first

known arrival of BCNH at Lake Calumet Wetlands (LCW) are based on periodic checks each spring of known previous nesting locations. Post-breeding surveys were initiated June 12 and concluded on August 21.

RESULTS

Nesting Habitat at Lake Calumet Wetlands 1984–2003

Throughout much of the early part of the 20th century, the Lake Calumet BCNH colony was located along the Calumet River just north of the confluence with the Grand Calumet River (Landing 1986). The construction of the Thomas J. O'Brien Lock and Dam resulted in the BCNH relocating to stand of eastern cottonwood (*Populus deltoides*) trees at Lake Calumet's Big Marsh (Fig. 1) during the early 1970s. The BCNH nested in the *Phragmites* at Big Marsh from 1984 through 1998; however, prolonged high-water levels in 1999 killed the *Phragmites* used for nesting structure. Although the emergent vegetation has recovered, as of 2003 the BCNH had not returned to nest at Big Marsh.

The cottonwoods at IRM were second in importance as a nesting site for this colony over the past 19 years (Table 1). The herons nested in the cottonwoods along the Calumet River at the southern portion of IRM during 1987–89; their abandonment of that rookery coincided with nesting by Red-tailed Hawks (*Buteo jamaicensis*) at that location in 1990. The BCNH began nesting at the cottonwood grove at IRM near 122nd St. in 1991, but have not nested at that site since 1996 after a pair of Red-tailed Hawks nested there. A lack of tree regeneration coupled with the death/poor condition of larger trees due to beaver (*Castor canadensis*) damage or high-water levels have been suggested as the causes for the failure of BCNH to return to nest in the cottonwoods. The *Phragmites* at the northern portion of IRM and at Heron Pond have become important nesting sites in recent years (Fig. 1).

During 2002 and 2003 the portion of IRM occupied by the nesting BCNH was characterized as a hemi-marsh condition, i.e., irregular-shaped and -sized stands of *Phragmites* interspersed with open water. The *Phragmites* cover ranged from narrow "fingers" and




Figure 1. Map of southeastern Chicago, Illinois, and northwestern Indiana, showing locations used by nesting Black-crowned Night-Herons.

Table 1. Known Black-crowned Night-Heron nesting locations at Lake Calumet Wetlands, 1984–2003.

Location	Year	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	
Indian Ridge Marsh (IRM) <i>Phragmites</i>											X	X					X	X		X	X	
IRM S. cottonwoods					X	X	X															
IRM N. cottonwoods									X	X	X	X	X									
Big Marsh <i>Phragmites</i>		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X						
Heron Pond <i>Phragmites</i>													X	X					X	X		

smaller isolated patches or clumps to larger blocks of cover. Thus, the area where the breeding colony was located was heterogenous with regard to available cover.

The BCNH colony was comprised of three "sub-colonies," each isolated from each other by an expanse of open water (Fig. 2). The northern subcolony held a greater proportion of the nests than either the eastern or southern subcolonies. Isolation of nesting areas from the shoreline by deeper water and thick emergent cover may have reduced the incidence of mammalian predation and vandalism/disturbance. Least isolated from shoreline by distance or deep, open water, nests in the eastern subcolony did experience greater predation loss in 2002 than in the monitored portion of northern subcolony.

The BCNH constructed nests of the previous year's *Phragmites* stems (occasionally lined with purple loosestrife (*Lythrum salicaria*) or unidentified woody stems) placed within a clump of standing stems. Nest distribution was generally clumped. There were concentrations of nests in close proximity (some less than a meter apart), generally in areas of sparser cover (i.e., near edges of cover or where *Phragmites* stems were less dense). Occasional nests were located in single clumps of reeds or scattered in denser cover.

Arrival and Courtship

The earliest date that BCNH have been observed at LCW during the period 1993 to 2003 was on March 10, 1994 (Table 2). During 1993, 1994, 1995, 2000, 2001, and 2003 no herons were observed on earlier (i.e., prior to date first observed) visits; no previous visits were recorded for 1996, 1997, 1998, and 2002. No BCNH were observed on March 29, 1992 or March 23, 1999, thus the herons apparently began arriving after those dates. Although we cannot be sure of the exact date of arrival based on these data, they do indicate that BCNH typically began arriving in substantial numbers during the latter half of March.

BCNH may continue to arrive at LCW well after others have begun nest building and pairing. In 2002 the number of BCNH at IRM greatly increased between April 26 and May 12. Also, no BCNH were observed at nearby Heron Pond (HP) until April 6 of that year. Similarly, BCNH were present at IRM as early as March 10 in 1994, but were not observed at

nearby Big Marsh (BM) as late as May 1, though were known to be successfully nesting there later in the season.

In 2002 BCNH were first observed at LCW on March 17, and were initially seen (n=4) carrying sticks on April 6. Early in the breeding cycle this behavior signifies beginning of nest building by males just prior to courtship and accepting a female at the nest (Meyerriecks 1960, Palmer 1962). Later, after pair formation, the male will present twigs to females prior to copulation and this behavior persists until after the eggs are laid. Gross (1923) reported that the first eggs were laid an average of seven days after start of nest construction; based on this information the first eggs would have appeared at LCW around April 13.

The first pairs were observed at nests at IRM on April 16; the number of pairs seen from an unobstructed vantage point increased from 2 on April 16, to 10 April 23, and 22 on May 2. Circling flights (synchronization of sexual behavior in early pair formation, Meyerriecks 1960) were observed on April 16; precopulatory displays (including billing, feather nibbling, twig presentation, erection of feathers, and neck stretch), and copulation were recorded as early as April 18 and April 19. According to Allen and Mangels (1940) the first eggs are laid an average of 3.3 days after copulation or 4–5 days after pair formation. Based on this information the first eggs would have appeared on April 20 or 21.

In 2003 BCNH were first observed carrying sticks on April 9, thus the first eggs would have appeared at LCW around April 16; in fact, pairs were first observed at nests on April 16. Precopulatory displays (including billing, feather nibbling, twig presentation, erection of feathers, and neck stretch) were recorded as early as April 18, thus the first eggs would be expected to have appeared on April 21. Based on our sample of nests, the first clutches were initiated on April 16, with half the nests initiated by May 8, and the last on May 23.

Nesting Phenology

The phenology of BCNH clutch initiation, egg hatching, and juvenile fledging/dispersal at IRM was reconstructed based on direct observation (i.e., nest monitoring) and the following assumptions:

BCNH Nesting Colonies (2002–2003)
Lake Calumet Area




Figure. 2. Location of Black-crowned Night-Heron colony in *Phragmites* cover at Indian Ridge Marsh during 2002 and 2003 and Heron Pond in 2002.

Table 2. Number at first sighting by week of occurrence of Black-crowned Night Herons at Lake Calumet Wetlands, 1993–2003.

Year	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Week											
March 1–7							No Data				
March 8–14		1									
March 15–21									16		
March 22–28			14		40	3		2		18	
March 29–April 4	25			17					75		

- 1) approximately two days to lay each egg (Gross 1923, Palmer 1962, Tremblay and Ellison 1980);
- 2) an average incubation period of 24 (23.5) days for A-eggs (i.e., first egg laid; Custer et al. 1992);
- 3) hatching one day after pipping, two days after cracking (Custer and Peterson 1991);
- 4) B-egg hatches about one day after A-, and C- hatches two days after B- (Custer et al. 1992); we assumed D-egg hatched one day after the C-egg;
- 5) flight is attained at about six weeks (42 days) of age (Palmer 1962, Wolford and Boag 1971);
- 6) on average, young of the year disperse at 58 days of age (Erwin et al. 1996).

In 2002 we initially marked 30 nests on May 15 and added new nests as encountered in that portion of the colony (n= 9 on May 20, n= 6 on May 26, n= 2 on May 30, and n= 1 on June 4). Based on our monitored sample of nests, we estimated that clutches were initiated as early as April 20, with half being initiated by mid-May, and the last on June 7 (Fig. 3). The last clutches initiated were in recycled nests; excluding second and third clutches (i.e., recycled nests) the latest a clutch was initiated, based on our marked sample, was May 23.

The estimate of April 20 for the initiation of laying coincides with estimates of clutch initiation based on our observations of pre-copulatory displays and copulation and information provided in Allen and Mangels (1940).

The first eggs would have hatched by mid-May, with A-eggs from half of the monitored nests hatching by the second week in June; A-eggs from the last nests initiated would have hatched on July 1 (Fig. 3). Thus, eggs would have been present at IRM over a period of about two and one-half months (April 20 to July 4-5).

Based on a flightless period of six weeks (Palmer 1962), the first and last young, respectively, attained flight on June 25 and August 10 (Fig. 3). Erwin et al. (1996) reported that age of dispersal in BCNH averaged 55 and 60 days in the two years of their study. Thus, based on an average dispersal age of 58 days, dispersal of juvenile IRM BCNH began about mid-July and continued through late August, with most having reached dispersal age by the middle of August. This coincides well with the results of our post-breeding surveys, which revealed a precipitous decline in juveniles observed at IRM between August 10 and 24 (see Fledging and Dispersal below).

In 2003 the first eggs would have hatched by mid-May; A-eggs from half of the monitored nests would have hatched by early June; A-eggs from the last nests initiated would have hatched on June 16 (Fig. 4). Thus, eggs would have been present at IRM over a period of about two




Figure 3. Nesting phenology of Lake Calumet Black-crowned Night-Herons, 2002.




Figure 4. Nesting phenology of Lake Calumet Black-crowned Night-Herons, 2003.

months. The first young attained flight on July 5, and dispersal of juvenile BCNH from IRM began the first week in July and continued through mid-August, with most having reached dispersal age by the end of July. This coincides well with the results of our post-breeding surveys which revealed that most juveniles had left by mid-August of that year (see Fledging and Dispersal below).

Clutch Size

Mean clutch size (all clutches) of BCNH nesting at IRM in 2002 was similar to that of clutches in tree nests at nearby ISPAT Inland Steel (IIS) (Table 3). Mean clutch sizes (all clutches) at IRM in 2002 and 2003 were very similar; clutch size declined considerably at IIS in 2003.

A total of six monitored nests at IRM was recycled (one nest with second and third clutches) in 2002. Mean clutch size for initial clutches was slightly higher than for all clutches (Table 3). We did not document any recycled nests at IRM in 2003.

When sorted by median date of initiation (5/13, see Nesting Phenology above), i.e., early nests < median date, late nests > median date, the average clutch size in 2002 was 0.48 eggs larger in nests initiated on or prior to the median date. Thus, early clutches were nearly

16% larger than late clutches; however, we did not detect this difference statistically ($t_{28} = 1.8, P = 0.08$).

At least four nests at IRM in 2002 had additional eggs laid in them after the original clutches were complete (based on the time interval between last egg of the clutch laid and presence of new eggs). In two cases one additional egg was laid after clutches of three were

complete; in the other two nests, one or more of the original eggs were found in the water and new eggs added to the nest. In one of these, a known total of five eggs were laid. In the other, the nest was recycled following the disappearance of the original clutch of three eggs. At least six eggs were subsequently found in the nest, with all eventually disappearing; four eggs (some from both clutches) were found in the water below this nest. We did not identify any such “dump nests” at IRM in 2003.

Nest, Egg, and Nesting Fate

The most common cause of failed clutches at IRM in 2002 was through poorly constructed nests, which allowed eggs to roll out into the water (Table 4). Predation was the next most common cause of nest failure; the complete disappearance of two clutches/broods may also have represented predation events. Including these, 7 of 51 (13.7%; includes second clutches) monitored clutches/broods were lost to predation at IRM. This amounted to the loss of 13 eggs and 7 chicks; coupled with 2 eggs with bill marks found in completed clutches, a minimum of 22 eggs or nestlings were lost to nest predators. It was difficult to determine the cause of death of most nestlings due to condition of carcasses upon discovery. BCNH

Location/Status	n	Mean	S.E.	Min-Max	Mode	Comments	Citation
IRM 2002 — all clutches 2002	40	3.40	0.11	2-5	3	includes second clutches; nearly bimodal (3- and 4-egg clutches)	this study
IRM 2002 — first clutches only	36	3.42	0.12	2-5	3/4	excluding second clutches; bimodal	this study
IRM 2002 — first clutches ≤ May 13	18	3.56	0.17	2-5	4	initiated prior to or on median date; nearly bimodal (3- and 4-egg clutches)	this study
IRM 2003 — all clutches		3.36					this study
IRM 2002- first clutches > May 13	17	3.08	0.19	2-4	3	initiated after median date	this study
ISPAT Inland Steel, IN 2002 — first clutches	13	3.38	0.21	3-5	3	tree nests; exposure to environmental contaminants likely	this study
ISPAT Inland Steel, IN 2003		3.06				tree nests; exposure to environmental contaminants likely	this study
Maryland	69	3.46	NA	2-5	4	nearly bimodal (3- & 4-egg clutches); shrub/tree nests; no apparent impact environmental contaminants	Rattner et al. 2001
Oregon and Washington — all clutches	260	3.45 ^{1,2}	0.08-0.11	2-5	NA	means 3.10 to 3.92; shrub/tree nests; no apparent impact environmental contaminants	Blus et al. 1997
Oregon and Washington — first clutches	208	3.47 ^{1,2}	0.08-0.12	2-5	NA	means 3.09 to 3.93; shrub/tree nests; no apparent impact environmental contaminants	Blus et al. 1997
Idaho	281	3.6 ^{1,2}	0.08-0.13	1-6	3/4 ²	means 3.4 to 3.7; shrub/trees; DDE impacts on reproduction	Findholt and Trost 1985
NV, OR, WA — ≤ 8 ppm DDE	133	3.75 ^{1,2,3}	0.15 ¹	NA	NA	means 3.67 to 3.80; shrub/tree nests; no apparent impacts environmental contamination	Henny et al. 1984
NV, OR, WA — > 8 ppm DDE	50	3.17 ^{1,2,3}	0.19 ¹	NA	NA	means 3.00 to 3.41; shrub/tree nests; DDE impacts on reproduction	Henny et al. 1984
Massachusetts and Rhode Island	346	3.79 ^{1,2,3}	NA	2-5	4	means 3.66 to 3.96; possible DDE impacts on reproduction	Custer et al. 1983
North Carolina	121	3.32 ^{1,2,3}	NA	2-5	3	means 3.08 to 3.50; no apparent impacts environmental contaminants	Custer et al. 1983
Quebec	98	4.1 ^{1,2,3}	0.1	1-6	NA	means 3.9 to 4.2; shrub/tree nests; no apparent impacts environmental contaminants	Tremblay and Ellison 1980
Alberta — first clutches	116	3.60 ^{1,2,3}	NA	1-6	3/4 ^{2,3}	means 3.2 and 4.0; emergent vegetation nests	Wolford and Boag 1971
CA — pre-1947	35	3.86	NA	NA	NA	unknown status	(see Henny et al. 1984)
NV, OR, WA — pre-1947	41	3.80 ^{1,2,3}	NA	2-5	NA	unknown status	Henny et al. 1984
Utah — pre-1947	41	4.1	0.13	2-6	4	unknown status	Findholt and Trost 1985

¹ Grand mean; ² Multiple colonies; ³ Multiple years

Fate of Clutch	Number of Clutches	Percent of Clutches
Successful ¹	34	66.7
Eggs Rolled Out, Poorly Constructed Nest	7	13.7
Depredated ²	5	9.8
Abandoned ³	3	5.9
Eggs/Nestlings Gone, No Signs	2	3.9

¹ at least one nestling survived to 15 days post hatch

² one clutch assumed abandoned after one egg depredated

³ one damaged egg abandoned

Notes-

a) another 4 broods known to completely fail after 15 days

b) determined minimum 6 predation events (clutch/brood or single eggs/chicks)

c) 11 chicks dead in nest= 3 depredated (same nest), 8 unknown fate

eggs hatch asynchronously and later-hatching nestlings may be disadvantaged, with competition by older siblings leading to starvation, drowning, or trampling of younger nestlings. This was undoubtedly the cause of the disappearance of many nestlings. We frequently visited nests with two or three thriving siblings, and the smaller, less thrifty individual was often missing at our next visit.

Several eggs had apparently been depredated by Ring-billed Gulls (RBG), judging by bill marks, although in at least one case the gulls may have scavenged a previously abandoned clutch. Large numbers of RBG frequented the area and were often seen fishing in open-water portions of IRM. We did not observe Gulls at BCNH nests, although we did note a BCNH briefly chase a RBG that was flying low over the colony. We did find and examine three siblings < 5 days old that had been killed by single pecks to the back of their skull.

Most nests successfully fledged at least one young on 2003 (Table 5); there were no known or suspected losses of entire clutches or broods to predation. Although no eggs were observed in water, all eggs missing from two flat-topped nests were assumed to have rolled out into the water. One nest was presumed abandoned as the marked eggs were present well after they should have hatched. At least 25 eggs were missing or otherwise failed to hatch. Twenty-three nestlings did not survive to fledging (i.e., 15 days post-hatch); of these, 21 were missing, 1 was found dead in nest (cause unknown), another had fallen out of the nest and was found with its neck caught between *Phragmites* stems.

Fledging and Dispersal

The number of juveniles observed in 2002 increased dramatically between mid-July and mid-August. The first appearance of juveniles between June 29 and July 13 coincides rather well with our first observation of flighted juveniles on July 3 while performing nest checks, and estimated dates that older chicks attained flight. The number of juveniles observed peaked on August 10, before declining precipitously between August 10 and 24 (Fig. 5). This reduction in juveniles is consistent with our expectation, based on nesting phenology (see above), that 97% of juveniles would have reached dispersal age (~58 days

post-hatch) by August 24. In 2003 the number of juveniles observed peaked earlier than in 2002; in 2003 most young were gone from IRM by mid-August (Fig. 5).

Productivity

While comparisons of productivity between studies are useful, the ratio of breeders to non-breeders, food availability, local weather patterns, climate/growing season (latitude), predation, and a variety of density-dependent factors can influence productivity. Most powerful are comparisons with contemporary local/regional populations. The only sizable colony in the south Chicago area available to us for comparison during 2002 and 2003 was at IIS, located on the Lake Michigan shoreline only 15 km from the Lake Calumet colony (Fig. 1). This colony is located at the same latitude as LCW; the IIS BCNH nested in cottonwood trees during this study.

Daily nest survival rate was similar between the colonies we examined (Table 6). In 2002 clutches at IRM and IIS had a lower likelihood of surviving the incubation period, when compared to 2003 when the survival rate of nests to hatch was relatively high at both locations.

Daily survival rate of nests during the nestling period was similar between the IRM and IIS colonies (Table 6). Survival of broods to 15 days of age was lower at IIS than at IRM; brood survival at IRM was similar between years.

In 2002, nest success (a function of survival rate to hatching and survival rate to fledging) at IRM was greater than at IIS, which was poor (Table 6). Nest success at both IRM and IIS improved between 2002 and 2003, although remained poor at the latter.

Egg success, the probability of an egg hatching, was low at IIS during 2002, though improved between 2002 and 2003 at this site and at IRM. Nestling survival was low at IIS in 2002; survival of young to 15 days improved at both IRM and IIS between years (Table 6).

The mean number of young surviving/nest was very low at IIS, whereas "recruitment" was relatively high at IRM by comparison (Table 6). Recruitment increased at both IRM and IIS between 2002 and 2003.

Fate of Clutch	Number of Clutches	Percent of Clutches
Successful ¹	52	94.5
Eggs Rolled Out, Poorly Constructed Nest	2	3.6
Depredated	0	0
Abandoned	1	1.8
Eggs/Nestlings Gone, No Signs	0	0

¹ at least one nestling survived to 15 days post-hatch

² no confirmed or suspected predation of marked nests

Notes-

a) 23 chicks failed to survive 15 days= 21 missing, 2 chicks dead in nest (1 accidental death, 1 unknown cause of mortality)




Figure 5. Numbers of juvenile Black-crowned Night-Herons observed at Indian Ridge Marsh colony during 2002 and 2003 post-breeding surveys.

Table 6. Reproductive parameters in a Black-crowned Night-Heron colony located in the Calumet region of Illinois and northwestern Indiana, 2002 and 2003.

Location	Indian Ridge Marsh—This Study 2002	Inland Steel —This Study 2002	Indian Ridge Marsh—This Study 2003	Inland Steel —This Study 2003	Maryland —Rattner et al. 2001	Pacific Northwest —Blus et al. 1997	San Francisco Bay —Hothem et al. 1995	East Coast —Custer et al. 1983
Nesting Parameters								
Egg Laying and Incubation (n)	46	15	55	17	69			531
Daily Survival Rate	0.9895	0.9855	0.9985	0.9931	0.9936	0.9240–0.9934 ⁴		
Survival Rate to Hatch	0.7762 ¹	0.7043 ¹	0.9635	0.8474	0.8517 ²		0.475–0.701	0.547–0.902 ²
Nesting Period (n)	37	13	52	15	59		395	(<531?)
Daily Survival Rate	0.9922	0.9864	0.9899	0.9809	0.9911	0.9782–1.000 ⁵		
Survival Rate to Fledging	0.8892 ³	0.6245 ³	0.8593	0.7485	0.8745 ³		0.691–0.896	0.758–1.000 ³
Nest Success	0.6920	0.4398	0.8280	0.6343	0.7448			0.530–0.867
Probability of Egg Hatching	0.8860	0.6852	0.9195	0.8333	0.8933			0.852–0.917
Probability Young Survived 15 days	0.8365	0.5143	0.8688	0.7714	0.8917			0.771–1.000
Egg Success	0.5115	0.1550	0.6614	0.4078	0.5932			0.385–0.716
Mean Clutch Size	3.40	3.38	3.36	3.12	3.46			3.08–3.96
Mean Number of Young Surviving 15 Days	1.74/nest	0.52/nest	2.22/nest	1.27/nest	2.05/nest	0.41–1.83/nest	1.45–2.04/nest	1.19–2.65

¹ = 24 day laying and incubation period; ² = 25 day laying and incubation period; ³ = 15 days post-hatch; ⁴ = 27 day laying and incubation period; ⁵ = 14 days post hatch

DISCUSSION

BCNH utilize a wide range of arboreal habitats for nesting including upland orchards (Gross 1923), bottomland forests (Bjorklund and Holm 1997), coastal islands (Parsons 1995), and trees in towns and cities (Farwell 1919, Cunningham 1945). Breeding colonies are also frequently located in emergent vegetation such as *Phragmites* or cattails (*Typha*), with nests located just inches above the water (Nelson 1876–1877, Rockwell 1910, Greenwood 1981). During the 20 years from 1984 to 2003 the Lake Calumet BCNH changed nesting locations as habitat conditions dictated, i.e., when nesting cover was impacted by fire, flooding, or succession (or lack thereof). The colony has maintained a high degree of fidelity to this area in spite of these changes. This supports Graber et al.'s (1978) assertion that the quality of the surrounding foraging habitat is a more important criterion for nesting than the particular habitat in which the nests are actually placed.

There appeared to be a clear preference for relatively open nest sites (i.e., of lower *Phragmites* stem density). A more open nest would seem to result in greater nest predation risk. However, 1) nests tended to be clumped in these more open areas and there may be higher survival of individual nests within groups, and 2) there may be greater inclusive fitness in that parents may be better able to avoid predators at some risk of clutch/brood predation.

Clutch size may be affected by a variety of factors linked to female condition and/or environmental quality (Winkler and Allen 1996). Mean clutch size in our study in 2002 was similar to that of recently-studied colonies in Maryland and the Pacific Northwest. The average clutch size at IRM was at the upper end of the range of average clutch sizes observed in colonies that were thought to be impacted by DDE, and below the range of clutch sizes observed for "clean" colonies in the United States and Canada. Interestingly, mean clutch size for first and second clutches in our study differed by the same amount as those in colonies in Oregon and Washington. The reason for the decline in clutch size between 2002 and 2003 at IIS was unknown.

Early clutches (< median date) were larger, as compared with those completed later (> median date). Custer et al. (1983) noted that BCNH clutches initiated later in the

season had a larger proportion of smaller clutches (i.e., < four eggs) than earlier clutches. It is unclear as to whether such seasonal declines in clutch size are related directly to female condition or an adaptive response to declining resources as nesting season progresses (Winkler and Allen 1996).

Our estimate of April 20 to June 10 for the egg-laying period at IRM in 2002 is similar to April 24 to June 18 reported by Graber et al. (1978) for northern Illinois colonies. However, our estimate of April 20 to May 26 for the egg-laying period in 2003 is somewhat shorter than reported by Graber et al. (1978). In 2002, eggs would have been present at IRM over a period of two and one-half months; this is a longer egg period than in 2003 by about two weeks. This is consistent with the number of juveniles present during our post-breeding surveys; most young were gone from IRM by mid-August of that year, or about two weeks earlier than in 2002. We did not encounter recycled nests in 2003, thus there may have been fewer re-nesting attempts in that year.

Daily survival rate of nests at IRM and IIS during the nestling period were similar to those examined by others. Survival of broods to 15 days of age was lower in tree-nesting BCNH at IIS than the *Phragmites*-nesting colony at IRM or reported in previous studies. Parsons (1995) reported that nestling survival was lower in Cattle Egrets nesting in shrubs as compared to those nesting in *Phragmites* in the same colony (Parsons 1995). We sometimes noted eggs and carcasses of young on the ground below nesting trees at IIS. This colony was directly exposed to winds off of Lake Michigan, and we assume that some of this loss may have been due to high winds and other mishaps causing nestlings to fall out of trees. Brood survival at IRM was similar between years, and was relatively high in comparison with past studies.

Although some eggs and young were lost to predation, we were not able to determine the cause of all losses. It is plausible that RBG were responsible for the disappearance of some of the eggs and smaller nestlings that weren't accounted for. However, predation by this species did not seem to be an important factor at IRM, particularly in light of the large numbers of RBG frequenting this wetland. In contrast, RBG were responsible for the poor productivity of some BCNH colonies in Alberta (Wolford and Boag 1971). Predation by Common Ravens (*Corvus corax*) essentially eliminated production in one colony in the

western United States (Henny et al. 1984), and egg depredation by American Crows (*Corvus brachyrhynchos*) affected nest success in some colonies in the Pacific Northwest (Blus et al. 1997). Although gulls, crows, and other avian predators were present, predation did not have a demonstrable impact on productivity of the LCW BCNH colony.

Henny (1972) determined that 2.0–2.1 young produced/breeding pair is needed to maintain BCNH populations, and Wolford and Boag (1971) calculated that a colony producing 1.1 fledged young/pair would disappear in 20 years. Thus, the estimated fledging rates (actually survival to 15 days) at IRM in 2002 in this study may not have been high enough to maintain a stable population. However, recruitment in 2003 was among the highest reported in other studies. It is not unusual for productivity of BCNH colonies to vary between years. For example, Greenwood (1981) reported annual production of 0.23, 0.57, and 2.20 fledglings/pair for a colony occupying a marsh in North Dakota; strong storms with hail and high winds were responsible for large losses of nests and young in two of those years. However, in the absence of such catastrophic events (including heavy losses to predation) productivity of BCNH does not vary this dramatically between years (e.g., see Custer et al. 1983).

Exposure to environmental contaminants has been associated with impaired reproduction (Custer et al. 1983, McEwen et al. 1984, Henny et al. 1984, Findholt and Trost 1985, Hoffman et al. 1993, Hothem et al. 1995) and teratogenic effects (Hoffman et al. 1993, Hothem et al. 1995) in BCNH. At the time of this writing we are examining selected environmental contaminants and biomarkers of exposure in pipping embryos collected from this colony in 2002; the results of this work are not addressed here. However, we did not observe any gross deformities in pipping embryos or nestlings and, as noted above, reproductive parameters for BCNH nesting at IRM were within normal limits for this species. Thus, it would appear that environmental contaminants are not having observable impacts on the productivity of this population.

LITERATURE CITED

- Allen, R.P., and F.P. Mangels. 1940. Studies of the nesting behavior of the Black-Crowned Night Heron. Proc. Linn. Soc. New York 50/51:1–28.
- Bjorklund, R.G., and D.J. Holm. 1997. Impact of flooding on Illinois River wading bird colonies. Transactions of the Illinois State Academy of Sciences 90(3 and 4):123–133.
- Blus, L.J., B.A. Rattner, M.J. Melancon, and C.J. Henny. 1997. Reproduction of Black-crowned Night-Herons related to predation and contaminants in Oregon and Washington, USA. Col. Waterbirds 20:185–197.
- Cunningham, J. W. 1945. Missouri region. Audubon Magazine 47(5):50–51.
- Custer, T.W., and D.W. Peterson, Jr. 1991. Growth rates of Great Egret, Snowy Egret, and Black-crowned Night-Heron chicks. Col. Waterbirds 14:46–50.
- Custer, T.W., G.L. Hensler, and T.E. Kaiser. 1983. Clutch size, reproductive success, and organochlorine contaminants in Atlantic coast Black-Crowned Night Herons. Auk 100:699–710.
- Custer, T.W., G.W. Pendleton, and R.W. Roach. 1992. Determination of hatching date for eggs of Black-crowned Night-Herons, Snowy Egrets, and Great Egrets. J. Field Ornithol. 63:145–154.
- Erwin, R.M., J.G. Haig, D.B. Stotts, and J.S. Hatfield. 1996. Reproductive success, growth, and survival of Black-Crowned Night Heron (*Nycticorax nycticorax*) and Snowy Egret (*Egretta thula*) chicks in coastal Virginia. Auk 113:119–130.
- Farwell, E.D. 1919. Bird observations near Chicago. Privately printed, Chicago, Illinois. 188 pp.
- Findholt, S.L. and C.H. Trost. 1985. Organochlorine pollutants, eggshell thickness, and reproductive success of Black-crowned Night-Herons in Idaho, 1979. Col. Waterbirds 8:32–41.
- Graber, J.W., R.G. Graber, and E.L. Kirk. 1978. Illinois birds: Ciconiiformes. IL Nat. Hist. Sur. Biol. Notes No. 19. 80 pp.


- Greenwood, R.J. 1981. Observations on Black-crowned Night-Heron breeding success in a North Dakota marsh. *Can. Field Nat.* 95:465–467.
- Gross, A.O. 1923. The Black-Crowned Night Heron (*Nycticorax nycticorax naevius*) of Sandy Neck. *Auk* 40:191–214.
- Henny, C.J. 1972. An analysis of the population dynamics of selected avian species with special reference to changes during the modern pesticide era. US Fish Wildl. Serv. Wildl. Res. Rpt. 1. 99 pp.
- Henny, C.J., L.J. Blus, A.J. Krynitsky, and C.M. Bunck. 1984. Current impact of DDE on Black-crowned Night-Herons in the intermountain west. *J. Wildl. Manage.* 48:1–13.
- Hoffman, D.J., G.J. Smith, and B.A. Rattner. 1993. Biomarkers of contaminant exposure in Common Terns and Black-Crowned Night-Herons in the Great Lakes. *Environ. Tox. Chem.*, 12:1095–1103.
- Hothem, R.L., D.L. Roster, K.A. King, T.J. Keldsen, K.C. Marois, and S.E. Wainwright. 1995. Spatial and temporal trends of contaminants in eggs of wading birds from San Francisco Bay, California. *Environ. Tox. Chem.* 14:131–1331.
- Landing, J.E. 1986. Conceptual plan for the Lake Calumet Ecological Park, Chicago, Illinois. 128 pp.
- Marcisz, W.J., J.M. Levensgood, A.M. Klement, and M.A. Kurcz. 2005. Population trends in a Black-crowned Night-Heron colony at Lake Calumet wetlands, Chicago, Illinois. *INHS Bulletin* 37(3).
- McEwen, L.C., C.J. Stafford, and G.L. Hensler. 1984. Organochlorine residues in eggs of Black-Crowned Night-Herons from Colorado and Wyoming. *Environ. Tox. Chem.* 3:367–376.
- Meyerriecks, A.J. 1960. Comparative breeding behavior of four species of North American herons. *Publ. Nuttall Ornithol. Club No. 2*, Nuttall Ornithol. Club, Cambridge, MA. 158 pp.
- Nelson, E. W. 1876–1877. Birds of north-eastern Illinois. *Essex Institute Bulletin* 8:9–12.
- Palmer, R.S. 1962. Handbook of North American birds. Volume 1 Loons through Flamingos. Yale Univ. Press, New Haven, CT. 567 pp.
- Parsons, K.C. 1995. Heron nesting at Pea Patch Island, Upper Delaware Bay USA: abundance and reproductive success. *Col. Waterbirds* 18:69–78.
- Rattner, B.A., P.C. McGowan, J.S. Hatfield, C.S. Hong, and S.G. Chu. 2001. Organochlorine contaminant exposure and reproductive success of Black-Crowned Night-Herons (*Nycticorax nycticorax*) nesting in Baltimore harbor, Maryland. *Arch. environ. Contam. Toxicol.* 41:73–82.
- Rockwell, R. B. 1910. Some Colorado night heron notes. *The Condor* 12:113, 121.
- Tremblay, J., and L.N. Ellison. 1980. Breeding success of the Black-crowned Night-Heron in the St. Lawrence Estuary. *Can. J. Zool.* 58:1259–1263.
- Winkler, D.W., and P.E. Allen. 1996. The seasonal decline in Tree Swallow clutch size: physiological constraint or strategic adjustment? *Ecology* 77:992–932.
- Wolford, J.W., and D.A. Boag. 1971. Distribution and biology of Black-crowned Night-Herons in Alberta. *Can. Field-Nat.* 85:13–19.

ACKNOWLEDGMENTS

We gratefully acknowledge the following for their assistance: Tom Barnett, ISPAT International; Robert Page, EarthTech; Marv Piwoni, Julie Hafermann, WMRC; Suzanne Malec and Nicole Kamins, City of Chicago Dept. of Environment (DOE); John Dassow, Loren Kirkwood, and Jyotsana Agarwal, INHS; Tom Custer, USGS; Dick Bjorklund, foraging survey volunteers, and three reviewers. This project was funded by the Illinois Waste Management and Research Center (Illinois Department of Natural Resources), City of Chicago Department of Environment, and the Illinois Wildlife Preservation Fund. The Illinois Department of Transportation—Aerial Surveys contributed services in-kind. Publication costs were partially funded by the Illinois Natural History Survey.

ILLINOIS
NATURAL
HISTORY
SURVEY

Black-crowned Night-Herons of the Lake Calumet Region, Chicago, Illinois



Article 4

Population Trends in a Black-crowned Night-Heron Colony at Lake Calumet Wetlands

Walter J. Marcisz, Jeffrey M. Levensgood,
Allison M. Klement, and Margaret A. Kurcz

Correspondence to:
Jeffrey M. Levensgood
Center for Wildlife and Plant Ecology
Illinois Natural History Survey
607 E. Peabody Dr.
Champaign IL 61820
217/333-6767
jleven@inhs.uiuc.edu

CONTENTS

ABSTRACT	111
INTRODUCTION	111
METHODS	111
Population Levels	111
Peak Numbers	111
Number of Nests	113
RESULTS	113
DISCUSSION	115
LITERATURE CITED	116
ACKNOWLEDGEMENTS	117

ABSTRACT

The number of active Black-crowned Night-Heron (*Nycticorax nycticorax*) nesting colonies in Illinois has declined significantly over the past century. Habitat loss/degradation and other factors such as exposure to environmental contaminants and competition for nest sites at established colonies may have contributed to this decline. In this study, we examined recent trends in population levels of Black-crowned Night-Herons nesting at wetlands associated with Lake Calumet in southeastern Chicago, Illinois. The number of Black-crowned Night-Herons nesting annually at these wetlands has fluctuated widely over the last two decades. Immigration of herons from riverine colonies may have driven population increases during the mid-1980s and early 1990s. However, this population has remained relatively stable at between 300 and 400 pairs during 1997–2003.

INTRODUCTION

The decline of the colonial-nesting Black-crowned Night-Heron (*Nycticorax nycticorax*) (BCNH) as a breeding species in the state of Illinois has been documented by a number of authors. According to Bohlen (1989:16), "The Black-crowned, like other heron species, is declining in numbers and has been for the past half-century." Mlodinow (1984:76) reported that, in the Chicago area, "[BCNH] Nesting colonies are steadily becoming smaller and fewer..." And Graber, et al. (1978:54) indicated that their Illinois data suggested "a serious decline in the BCNH population in the past half-century." Results of the Illinois Colonial Waterbird Survey indicated that there were an estimated 1,900 nesting pairs in 1987 (Kleen 1987), compared with 400+ in 1999 (Kleen 1999).

Young-of-the-year BCNH were reported in the Calumet marshes as early as 1874 (Nelson 1876–1877). Although there are no accurate figures for the extent of these wetlands during presettlement times, historic maps suggest that marshland formations totaled at least 22,000 acres on the Illinois side of the state line. Due to the impacts of industry, railroads, waste disposal, urbanization, and hydrologic changes during the 20th century, only approximately 500 acres of wetlands remained at Lake Calumet (LCW) in recent

years (Landing 1986), much of it in an impaired state. Despite these habitat losses and degradation, BCNH have maintained a tradition of nesting at wetlands associated with LCW in southeastern Chicago (Fig.1). The common reed (*Phragmites australis*) has been of primary importance as nesting cover for this colony since 1984; cottonwood trees (*Populus deltoides*) were of secondary importance in some years.

We present the results of weekly population censuses conducted during spring of 2002 and 2003 of BCNH nesting at LCW, along with a compilation of the results of censuses of nests and/or breeding BCNH conducted since 1984.

METHODS

Population Levels—BCNH are largely crepuscular/nocturnal foragers that disperse from their daytime roosts/nesting areas at dusk. During the 2002 and 2003 breeding seasons, BCNH population censuses were conducted at dusk at the northern portion of Indian Ridge Marsh on a weekly basis from the arrival of the BCNH in late March through late May (the final census in 2003 was conducted on June 1). Censuses were conducted under conditions of no precipitation and winds < 15 mph by two observers beginning one hour before sunset and continuing until the rate of departure was < 2 herons/5 minute period. As days lengthened the herons became active earlier and as of May 1 censuses were initiated two hours before sunset.

Although no BCNH were noted at nearby Heron Pond on March 24 and 30, 2002, a number of herons were observed at this site on April 6, 2002. After mid-April 2002 it became apparent that some herons were nesting at Heron Pond and censuses were subsequently conducted on April 26 and May 12, 2002 at that site. Early in the spring of 2003, a wildfire at Heron Pond destroyed essentially all of the emergent vegetation at that site, eliminating the BCNH's nesting substrate for the season. No BCNH nesting occurred at Heron Pond that year.

Peak Numbers—Peak numbers of BCNH were based on evening censuses of BCNH leaving nesting colonies to forage, conducted during 1992–2003. Data for 1992–2001




Figure 1. Map of southeastern Chicago, Illinois, and northwestern Indiana, showing locations used by nesting Black-crowned Night-Herons.

represented an average of 2 to 3 counts conducted during late-April/early-May of each year; the 2002 data represented the maximum of 11 counts conducted March 24 to May 26; the 2003 data represented the maximum of 11 counts conducted from March 23 to June 1. These were combined with counts of occupied tree nests (multiplied by 2 BCNH) conducted during 1987–89 and 1991–95.

Number of Nests—The number of nests for each year were calculated from actual nest counts conducted during 1984–91, combined with estimates of nesting pairs for the years 1992–2003. We compiled the results of annual nest counts conducted at Big Marsh during mid- to late-May of 1984–91 (no data for 1990). These counts consisted of a line of five to seven observers walking transects through the *Phragmites* stands and counting nests seen to one side of each person. A portion of the colony was not surveyed in 1986 and 1987; it was estimated that <15% of the nesting colony was missed, thus we added 15% to the nest counts in those years. Direct counts of tree nests located at Indian Ridge Marsh during 1987–91 were added to these counts. The number of nests present during 1992–2003 were estimated by dividing peak counts of BCNH by two to represent a pair for each nest. We feel that this provides a reasonably accurate approximation of breeding pairs (nests) as: 1) our data suggests that prior to incubation the vast majority of BCNH are counted during evening censuses, 2) 99.3% of known-aged herons observed during our evening population censuses were >1 year old, and 3) there are numerous accounts of nesting by one-year-old BCNH (Gross 1923, Nobel and Wurm 1942, Custer and Davis 1982). In 1994 the herons began nesting at nearby Big Marsh (in addition to the *Phragmites* and cottonwoods at Indian Ridge Marsh) late in the season. Thus, a complete count was not conducted at that location. Consequently, the 1994 estimate of 410 nests represents a minimum and was not included in Figure 4.

RESULTS

In 2002, BCNH were first noted at LCW on March 17; numbers gradually increased from the start of the evening population censuses on March 24 through May 12, when they peaked

at 750 herons (Fig. 2). The number of BCNH observed during evening censuses declined dramatically between May 12 and May 26; counts were discontinued after that time. This decline in BCNH leaving Indian Ridge Marsh during evening hours was undoubtedly due to the increasing numbers of herons remaining at nests to incubate eggs or brood young. Also, counting became more difficult as foraging trips became shorter in duration as BCNH returned to the colony to feed young or allow their mates to recess. No BCNH were observed at Heron Pond on March 24 or 30; however, 33 BCNH were counted there on April 6. Population censuses conducted on evenings of April 29 and May 10 indicated that a substantial number of BCNH were nesting there (142 counted on May 10).

BCNH were first noted at LCW on March 22 of 2003; with the exception of a leveling off of numbers between the weeks of March 30 and April 11, numbers gradually increased from the start of the evening population censuses on March 23 through May 10, when they peaked at 641 herons (Fig. 2). The number of BCNH observed during evening censuses declined dramatically between May 10 and June 1; evening censuses were discontinued after that time. As in 2002, this decline in BCNH observed was undoubtedly due to increasing numbers of uncounted incubating/brooding herons coupled with shorter, more frequent foraging trips by observed herons. Thus, the 2003 peak count of 641 BCNH was down 15% from the 2002 peak count of 750, but the estimated number of nests ($641 \text{ total BCNH} / 2 \text{ BCNH/nest} = 320 \text{ nests}$) was within “normal” limits for the period 1997–2002 (303 to 404 nests).

The peak BCNH population at LCW varied considerably during 1992–97, from a high of nearly 1,600 in 1992 to less than 600 in 2000 (Fig. 3). The quadratic trend was significant and described a relatively large proportion of the variation in peak numbers. Although there was a declining population trend during much of the 1990s, numbers have more or less stabilized at 600–800 herons in recent years.

The number of nests varied considerably during 1984–97, from a low of 266 in 1984 to a high of 871 in 1992 (Fig. 4). The cubic trend was significant and described a relatively large portion of the variation in number of nests. The number of BCNH nests increased dramati-




Figure 2. Results of counts of Black-crowned Night-Herons leaving nesting locations to forage during 2002 and 2003.




Figure 3. Peak numbers of Black-crowned Night-Herons at Lake Calumet wetlands, 1992–2003.

cally during the latter half of the 1980s and remained relatively high into the mid-1990s. Numbers of BCNH nests at LCW declined throughout much of the 1990s before essentially leveling off at 300–400 nests during the late 1990s and early 2000s (Fig. 4).

DISCUSSION

Many factors have been suggested as causes for recent declines in BCNH populations, including vandalism, disturbance at breeding colonies, drainage of wetlands, and land development for homes and recreation (Davis 1993). A relationship has been demonstrated between chlorinated hydrocarbon pesticide pollution and egg-shell thinning in herons (Faber et al. 1972). Pre-1947 (pre-DDT) and post-1947 (DDT-era) BCNH eggshell thickness measurements taken in 15 U.S. states demonstrate significantly thinner eggshells in post-1947 clutches (Ohlendorf and Marois 1990, Ohlendorf et al. 1978, Ohlendorf et al. 1977), but convincing documentation that DDT (DDE) and other pesticides have caused local BCNH population declines before DDT was banned in 1972 is lacking (Davis 1993).

High-water levels, resulting in flooded nests, have forced BCNH to relocate within LCW on numerous occasions over the past two decades. In virtually all of these cases, the BCNH completed a successful breeding season. In the Illinois River Valley of central Illinois, flooding has had an impact on heron colonies in recent years, with high water appearing to disadvantage BCNH in particular (Bjorklund and Holm 1997). The frequent and rapid flood peaks along the Illinois River have occurred as a result of floodplain constriction by levees, channelization of tributary streams, and other human activities (Bjorklund and Holm 1997).

Bjorklund and Holm (1997) found that prolonged elevated water levels at Illinois River sites during the period of nestling development (June through August) had a negative impact on the size of heron nesting populations during the following year's breeding season. Periods of major flooding along the Illinois River occurred during 1993–96, with corresponding decreases in nesting BCNH populations during that period (Bjorklund and Holm 1997). A spike in BCNH peak numbers/nests at LCW during the early




Figure 4. Number of Black-crowned Night-Heron nests at Lake Calumet wetlands, 1985–2003.

1990s corresponded to the severe flooding of the Illinois River floodplain (and resultant desertion of colonies by BCNH) during the same period, suggesting that emigration of BCNH from Illinois River colonies to LCW may have occurred during that time. The Breeding Bird Survey Trend Map (1966–1996) for BCNH also shows decreases ($< -1.5\%$) in Illinois River BCNH populations, and corresponding increases ($> +1.5\%$) in LCW BCNH populations during that period.

Interspecific competition for nest sites may have affected BCNH populations at various Illinois colonies. At Lake Renwick (Plainfield, Will Co., IL) Double-crested Cormorants (*Phalacrocorax auritus*) appear to have displaced Great Blue Herons (*Ardea herodias*), Great Egrets (*Ardea alba*), and BCNH from their traditional nest sites. As Double-crested Cormorant numbers increased, the BCNH population at Lake Renwick declined from 273 nests in 1983 to 57 nests in 1990 (Milosevich 1990). Numbers of Double-crested Cormorants have also increased at the Baker's Lake colony in northwestern Cook Co., while the BCNH population there declined from 220 nests in 1989 to 11 nests in 1992. It is plausible that BCNH emigrating from Lake Renwick, Baker's Lake, and other declining Illinois BCNH colonies contributed to the BCNH population increases at LCW during the late 1980s to early 1990s. The decline in the breeding population during the 1990s and relatively stable number of breeding pairs since that time suggests that LCW were ultimately unable to support a higher breeding population resulting from the emigration of BCNH from Illinois River and other colonies.

LITERATURE CITED

- Bjorklund, R.G., and D.J. Holm. 1997. Impact of flooding on Illinois River wading bird colonies. *Transactions of the Illinois State Academy of Sciences* 90(3 and 4):123–133.
- Bohlen, H.D. 1989. *The birds of Illinois*. Indiana University Press, Bloomington, Indiana. 222 pp.
- Custer, T.W., and W.E. Davis, Jr. 1982. Nesting by one-year-old Black-crowned Night-Herons on Hope Island, Rhode Island. *Auk* 99:784–786.
- Davis, W.E., Jr. 1993. Black-crowned Night-Heron (*Nycticorax nycticorax*). In *The Birds of North America*, No. 74 (A. Poole and F. Gill, eds.). Philadelphia: The Academy of Natural Sciences; Washington, D.C.: The American Ornithologists' Union.
- Faber, R.A., R.W. Risebrough, and H.M. Pratt. 1972. Organochlorines and mercury in common egrets and great blue herons. *Environmental Pollution* 3(2):111–122.
- Graber, J.W., R.R. Graber, and E.L. Kirk. 1978. Illinois birds: Ciconiiformes. *Biological Notes* No. 109. Illinois Natural History Survey, Champaign, Illinois. 80 pp.
- Gross, A.O. 1923. The Black-crowned Night-Heron (*Nycticorax nycticorax naevius*) of Sandy Neck. *Auk* 40:1–30; 191–214.
- Kleen, V.L. 1987. Illinois heron colony surveys—1987 report. *Illinois Birds and Birding* 3 (4):79–82.
- Kleen, V.L. 1999. 1999 heron colony survey report. Illinois Department of Natural Resources.
- Landing, J.E. 1986. Conceptual plan for the Lake Calumet Ecological Park: Chicago, Illinois. 128 pp.
- Milosevich, J.B. 1990. Lake Renwick—from gravel quarry to nature preserve. *Illinois Audubon* No. 233:2–6.

Mlodinow, S. 1984. Chicago area birds. Chicago Review Press, Chicago, Illinois. 220 pp.

Nelson, E.W. 1876–1877. Birds of north-eastern Illinois. Essex Institute Bulletin 8:9–12.

Noble, G.K., and M. Wurm. 1942. Further analysis of the social behavior of the Black-crowned Night-Heron. *Auk* 59:205–224.

Ohlendorf, H.M., and K.C. Marois. 1990. Organochlorines and selenium in California night-heron and egret eggs. *Environ. Monit. Assess.* 15(1):91–104.

Ohlendorf, H.M., E.E. Klaas, and T.E. Kaiser. 1977. Organochlorine residues and eggshell thinning in anhingas and waders. *Proc. Colon. Waterbird Group 1977*:185–195.

Ohlendorf, H.M., E.E. Klaas, and T.E. Kaiser. 1978. Environmental pollutants and eggshell thinning in the Black-crowned Night Heron. Pages 63–82 *in* *Wading Birds* (Sprunt, et al., eds.). Nat. Aud. Soc. Res. Rep. No. 7, New York.

ACKNOWLEDGMENTS

Funding for this study was provided by the Illinois Waste Management and Research Center of the Illinois Department of Natural Resources, the City of Chicago Department of Environment, and the Illinois Wildlife Preservation Fund. We thank Sue Elston, USEPA, for providing information on nest surveys conducted 1984–91, and Loren Kirkwood, INHS, for creating the map used in Figure 1. Publication costs were partially funded by the Illinois Natural History Survey.

NOTES